


Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



United States
Department of
Agriculture

Forest Service

Intermountain
Research Station

Research Note
INT-365

January 1987



HIDE2: Evaluation of Elk Hiding Cover Using a Personal Computer

L. Jack Lyon¹

ABSTRACT

An IBM² compatible program for personal computers is described. Input requirements are forest stand structural data; tree densities by diameter class. Output is an estimate of elk hiding cover values. Canopy diameter of shrubs and of trees with lower level foliage can be included in the evaluation. The program assumes random distribution of plants and produces a visual image representing 1 acre.

KEYWORDS: PC program, wildlife management, elk habitat

Coordination of elk habitat management and timber management on the National Forests requires comparison of existing elk habitat to habitat quality standards described in forest plans. These standards call for compilation of hiding cover and thermal cover values of forest stands. Identification of thermal cover, defined by crown canopy closure, poses no insurmountable problems because crown closure can be estimated from aerial photographs. Identification of hiding cover, however, can be extremely difficult because determination from aerial photographs is not possible, and there is rarely adequate time to obtain field samples.

This Research Note describes a program for personal computers that utilizes available timber stand information, produces hiding cover estimates with speed and precision, and runs on any IBM compatible PC (Lyon 1985). A mainframe version in FORTRAN is also available.

HIDE2: HOW IT WORKS

The currently accepted and widely used definition assumes that Hiding Cover is provided when vegetation hides 90 percent of an elk at 200 feet (Thomas and others 1979). Thus, in order to satisfy the definition, vegetation must visually obstruct the line of sight between an observer and an elk at a distance of 200 feet—and must do so in a block of sufficient width to hide an elk.

The PC program HIDE2 assumes an observer in a forest stand evaluating all points on an arc 200 feet away in increments of 1 inch. The percentage of such points made invisible by either stems or vegetation canopy is considered visual blockage. Visual blockage along the arc is evaluated in units 65 inches wide. By definition, only those units that provide 90 percent or greater visual blockage are classified as hiding cover.

The unique specialty of the PC program is that it displays the forest stand being evaluated on the screen. Input of already available stand inventory information, diameter (d.b.h.), and density (plants per acre) is requested sequentially. As answers are supplied, the following assumptions are made within the program:

Trees—all diameters up to 30 inches are considered to be tree stems. Visual blockage is created only by the stem.

Small Trees—diameters under 6 inches can be entered either as stems or as open-grown trees. Visual blockage by foliage is assumed when the diameter entry is followed by "+". The applicable foliage crown width is assumed to be 1 foot for each inch of stem diameter.

Shrubs—diameters from 30 to 90 inches are considered to be shrubs and are evaluated for visual blockage created by foliage.

As each class of vegetation is entered, random locations are generated for individual plants and the

¹Wildlife research biologist stationed at Intermountain Station's Forestry Sciences Laboratory, Missoula, MT.

²The use of trade or firm names in this publication is for reader information and does not imply endorsement by the U.S. Department of Agriculture of any product or service.

stand is plotted on the screen at about 1:500 scale for an area representing approximately 1 acre. Symbols for trees and shrubs of various types are appropriately scaled. Intermediate output consists of accumulated totals for Plant Density, Visual Blockage, and Hiding Cover.

Entry of a diameter greater than 90 terminates the program and produces summary output. In addition to Hiding Cover, output includes a 10-class frequency distribution of visual obstruction percentages.

HIDE2: DISTRIBUTION

The program HIDE2 is available from the author at no cost on receipt of a floppy disk. The program was written on an ATT PC6300, using MS-DOS 2.11 and GWBASIC 2.0. It has been tested for compatibility on an IBM PC, MS-DOS 2.10, and BASICA. It was designed for use with a color monitor, but it will run without modification on a monochrome monitor.

REFERENCES

Lyon, L. Jack. 1985. HIDE--modeling hiding cover for wildlife. PC program. In: Session III - Demonstrations of micro-computer applications: national workshop on micro-computer applications in fish and wildlife programs; 1985 October 9-11; Fort Collins, CO. Fort Collins, CO: Colorado State University. [Unnumbered].

Lyon, L. Jack; Marcum, C. Les. 1986. Field test of a PC program to evaluate hiding cover for elk. In: Proceedings, 1986 western states and provinces elk workshop; 1986 March 17-19; Coos Bay, OR. Portland, OR: Oregon Department of Fish and Wildlife: 271-274.

Thomas, Jack Ward; Black, Hugh, Jr.; Scherzinger, Richard J.; Pedersen, Richard J. 1979. Deer and elk. In: Wildlife habitats in managed forests: the Blue Mountains of Oregon and Washington. Agriculture Handbook 553. Washington, DC: U.S. Department of Agriculture, Forest Service: 104-127.

INTERMOUNTAIN RESEARCH STATION

The Intermountain Research Station provides scientific knowledge and technology to improve management, protection, and use of the forests and rangelands of the Intermountain West. Research is designed to meet the needs of National Forest managers, Federal and State agencies, industry, academic institutions, public and private organizations, and individuals. Results of research are made available through publications, symposia, workshops, training sessions, and personal contacts.

The Intermountain Research Station territory includes Montana, Idaho, Utah, Nevada, and western Wyoming. Eighty-five percent of the lands in the Station area, about 231 million acres, are classified as forest or rangeland. They include grasslands, deserts, shrublands, alpine areas, and forests. They provide fiber for forest industries, minerals and fossil fuels for energy and industrial development, water for domestic and industrial consumption, forage for livestock and wildlife, and recreation opportunities for millions of visitors.

Several Station units conduct research in additional western States, or have missions that are national or international in scope.

Station laboratories are located in:

Boise, Idaho

Bozeman, Montana (in cooperation with Montana State University)

Logan, Utah (in cooperation with Utah State University)

Missoula, Montana (in cooperation with the University of Montana)

Moscow, Idaho (in cooperation with the University of Idaho)

Ogden, Utah

Provo, Utah (in cooperation with Brigham Young University)

Reno, Nevada (in cooperation with the University of Nevada)